

LYMPHOCELES POST OPERATOIRES APRES LYMPHADENECTOMIES POUR CANCERS GYNECOLOGIQUES : TECHNIQUES DE PREVENTION ET PERSPECTIVES.

Aziz Achouri, Cyrille Huchon, Anne-Sophie Bats, Chérzade Bensaïd, Laurent Makké, Claude Nos, Fabrice Lécuru.

Service de Chirurgie Cancérologique Gynécologique et du Sein. Hôpital Européen Georges-Pompidou, Assistance Publique-Hôpitaux de Paris, Paris.

Université Paris Descartes, Sorbonne Paris Cité, Faculté de Médecine, Paris.

Correspondance : fabrice.lecuru@egp.aphp.fr

Bibliographie

1. Belot A, Grosclaude P, Bossard N, et al. Cancer incidence and mortality in France over the period 1980-2005. *Rev Epidemiol Sante Publique* 2008;56(3):159-75.
2. Selman TJ, Mann CH, Zamora J, Khan Kh S. A systematic review of tests for lymph node status in primary endometrial cancer. *BMC Women's Health* 2008;8:8.
3. Hawighorst H, Schoenberg SO, Knapstein PG, Knopp MV, Schaeffer U, Essig M. Staging of invasive cervical carcinoma and of pelvic lymph nodes by high resolution MRI with a phases array coil: comparison with pathological finding. *J Comput Assist Tomogr* 1998;22:75-81.
4. Selman JT, Mann C, Zamora J, et al. Diagnostic accuracy of tests for lymph node status in primary cervical cancer: a systematic review and metaanalysis. *CMAJ* 2008;178(7):855-62.
5. Tam KF, Lam KW, Chan KK, Ngan HYS. Natural history of pelvic lymphocysts as observed by. *Ultrasound Obstet Gynecol* 2008;32:87-90.
6. Scholz HS, Petru E, Benedicic C, Haas J, Tamussino K, Winter R. Fibrin Application for Preventing Lymphocysts after Retroperitoneal Lymphadenectomy in Patients with Gynecologic Malignancies. *Gynecologic Oncology* 2002;84:43-6.
7. Logmans A, Kruyt RH, de Bruin HG, Cox PH, Pillay M. Lymphedema and lymphocysts following lymphadenectomy may be prevented by omentoplasty: A pilot study. *Gynecol Oncol* 1999;75:323-7.
8. Benedetti-Panici P, Maneschi F, Buttillo G, et al. A randomized study comparing retroperitoneal drainage with no drainage after lymphadenectomy in gynecologic malignancies. *Gynecologic Oncology* 1997;65:478-82.
9. Suzuki M, Ohwada M, Sato I. Pelvic lymphocysts following retroperitoneal lymphadenectomy: retroperitoneal partial "no-closure" for ovarian and endometrial cancers. *J Surg Oncol* 1998;68(3):149-52.
10. Benedetti-Panici P, Maneschi F, Cuttillo G. Pelvic and aortic lymphadenectomy. *Surg Clin North Am* 2001 ;81:841-58.
11. Thomé Saint Paul M, Brémond A, Rochet Y. Absence de péritonisation après la chirurgie pelvienne carcinologique. *J Gynecol Obstet Biol Reprod* 1991;20:957-60.
12. Franchi M, Trimpos J, Zanaboni F, et al. Randomised trial of drains versus no drains following radical hysterectomy and pelvic lymph node dissection: a European Organisation for

Research and Treatment of Cancer Gynaecological Cancer Group (EORTC-GCG) study in 234 patients. *European Journal of Cancer* 2007;43:1265-8.

13. Charoenkwan K, Kietpeerakool C. Retroperitoneal drainage versus no drainage after pelvic lymphadenectomy for the prevention of lymphocyst formation in patients with gynaecological malignancies (Review). *The Cochrane Library* 2010;1.

14. Fujiwara K, Kigawa J, Hasegawa K, Nishimura R, Umezaki R, Anado M. Effect of simple omentoplasty and omentopexy in the prevention of complications after pelvic lymphadenectomy. *Int J Gynecol Cancer* 2003;13:61-6.

15. Benoit L, Boichot C, Cheynel N, et al. Preventing lymphedema and morbidity with an omentum flap after ilioinguinal lymph node dissection. *Ann Surg Oncol* 2005;12:793-9.

16. Shimotsuma M, Hagiwara A, Takahashi T, et al. Surface structure and cell zonation in human omental milky spots. *Lymphology* 1990;23:207-8.

17. Yamamoto R, Saitoh T, Kusaka T, et al. Prevention of lymphocyst formation following systematic lymphadenectomy. *Jpn J Clin Oncol* 2000;30:397-400.

18. Nezhat F, Yadav J, Rahaman J, Gretz H 3rd, Gardner GJ, Cohen CJ. Laparoscopic lymphadenectomy for gynecologic malignancies using ultrasonically activated shears: analysis of first 100 cases. *Gynecol Oncol* 2005;97(3):813-9.

19. Gallotta V, Fanfani F, Rossitto C, Vizzielli G et al. A randomized study comparing the use of the Ligaclip with bipolar energy to prevent lymphocele during laparoscopic pelvic lymphadenectomy for gynecologic cancer. *Am J Obstet Gynecol* 2010;203(5):483.e1-6.

20. Carcofaro P, Soliani G, Maestroni U, et al. Octreotide in the treatment of lymphorrhea after axillary node dissection: a prospective randomised controlled trial. *J Am Coll Surg* 2003;196:365-9.

21. Mahmoud S, Abdel-Elah K, Eldesoky AH, et al. Octreotide can control lymphorrhea after axillary node dissection in mastectomy operations. *Breast Journal* 2007;13(1):108-9.

22. Kim WT, Ham WS, KOO KC, Choi YD. Efficacy of octreotide for management of lymphorrhea after pelvic lymph node dissection in radical prostatectomy. *Urology* 2010;76(2):398-401.